

SHOCK

Dr. dr. Meddy Setiawan, SpPD, FINASIM

**FAKULTAS KEDOKTERAN
UNIVERSITAS MUHAMMADIYAH MALANG**



“Shock”

- Shock adalah suatu sindroma klinis yg ditandai dg kegagalan sistem sirkulasi untuk mempertahankan perfusi yg adekuat ke organ-organ vital tubuh

Klasifikasi Shock

(Menurut Hinshaw and Cox classification)



Hypovolemic	
Cardiogenic	<ul style="list-style-type: none">• Intrinsic• Kompresive
Distributive	<ul style="list-style-type: none">• Septic• Anaphylactic• Neurogenic
Obstructive	

Hypovolemic

Cardiogenic

- Intrinsic
- Kompresive

Distributive

- Septic
- Anaphylactic
- Neurogenic

Obstructive

Definisi

Hypovolemic shock

- Syok akibat volume darah dalam pembuluh darah yang kurang.

Cardiogenic shock

- Syok yg disebabkan kegagalan primer jantung dalam memompa. (pada keadaan volume intravaskuler yg cukup)


Distributive

- Syok akibat volume darah secara abnormal berkumpul dlm pembuluh darah perifer. Tjd gangguan distributive aliran darah,

Obstructive


- Gangguan kontraksi jantung akibat di luar jantung

Normal




Shock


Hypovolemic



Obstructive



Distributive



Etiologi

Hypovolemic Shock

- **Non hemmorrhagic:** vomiting, diarrhea, burns, environmental (dehydration)
- **Hemorrhagic**

Cardiogenic Shock

- Penyebab primer : miokarditis, peny jantung bawaan
- Penyebab sekunder : disfungsi miokard krn toksin, iskhemia


Distributive Shock

- Septic shock, SIRS, Anaphylaxis, Neurogenic shock, Insufficiency adrenal

Obstructive Shock

- Tamponade perikardiak
- Perikarditis restriktif
- Emboli paru (masif)
- Koarktasio aorta

...Effects of inadequate perfusion on cell function...





DEVELOPMENT OF SHOCK OR CIRCULATORY INSUFFICIENCY

EFFECTS

Reduced venous return
 ↓
 Reduced stroke volume
 ↓
 Reduced arterial pressure
 ↓
 Arteriolar- & veno-constriction
 ↓
 Reduced bloodflow
 ↓
 Hypoxia
 ↓
 Metabolic acidosis
 ↓
 Release of adrenaline
 " vasopressin
 " renin
 " red cells

EFFECTORS


 Baroreceptors & heart

 Vessels
 Cardiac-cerebral-hepatic-renal failure
 Tissue depression
 Chemoreceptors.

 Vessels, Kidney
 Spleen & blood

COMPENSATORY REACTIONS

Starling-law-reactions
 Starling-law-reactions
 Increased HR & contractility
 Reduced capillary pressure
 Hypoxia & glycolysis
 Renal tubular necrosis
 Increased ventilation
 Vasoconstriction.
 Water retention
 Vasoconstriction, Salt reabsorption
 Improved O₂-transport





Fig. 12-6





Table 264-2 Normal Hemodynamic Parameters

Parameter	Calculation	Normal Values
Cardiac output (CO)	SV × HR	4–8 L/min
Cardiac index (CI)	CO/BSA	2.6–4.2 (L/min)/m ²
Stroke volume (SV)	CO/HR	50–100 mL/beat
Systemic vascular resistance (SVR)	$[(MAP - RAP)/CO] \times 80$	700–1600 dynes · s/cm ⁵
Pulmonary vascular resistance (PVR)	$[(PAP_m - PCWP)/CO] \times 80$	20–130 dynes · s/cm ⁵
Left ventricular stroke work (LVSW)	SV(MAP - PCWP) × 0.0136	60–80 g-m/beat
Right ventricular stroke work (RVSW)	SV(PAP _m - RAP)	10–15 g-m/beat

...Immuno-inflammatory respon to shock...




Source: Fauci AS, Kasper DL, Braunwald E, Hauser SL, Longo DL, Jameson JL, Loscalzo J: *Harrison's Principles of Internal Medicine*, 17th Edition: <http://www.accessmedicine.com>

Copyright © The McGraw-Hill Companies, Inc. All rights reserved.


A schematic of the host immunoinflammatory response to shock.


Hypovolemic shock...




- **Mild (< 20% blood volume)**
 - Cool extremities
 - Increased capillary refill time
 - Diaphoresis
 - Collapsed veins
 - Anxiety
- **Moderate (20-40% blood volume)**
 - Tachycardia
 - Tachypnea
 - Oligouria
 - Postural changes
- **Severe (> 40% blood volume)**
 - Hemodynamic instability
 - Marked tachycardia
 - Hypotension
 - Coma

Cardiogenic shock...






Anaphylactic shock...



Neurogenic shock...



- As with hypovolemic shock but in high spinal injuries may also be accompanied by profound bradycardia due to loss of the cardiac accelerating nerve fibres from the sympathetic nervous system at T1-T4.
- The skin is warm and dry or a clear sweat line exists, above which the skin is diaphoretic.
- Priapism due to Peripheral nervous system stimulation

Septic shock...

- Similar to hypovolemic shock except in the first stages: Pyrexia (fever), due to increased level of cytokines
- Systemic vasodilation resulting in hypotension (low blood pressure)
- Warm and sweaty skin due to vasodilation
- Systemic leukocyte adhesion to endothelial tissue
- Reduced contractility of the heart
- Activation of the coagulation pathways, resulting in disseminated intravascular coagulation
- Increased levels of neutrophils

....MOGA BERMANFAAT...

